

E-ISSN: 2708-0021
 P-ISSN: 2708-0013
www.actajournal.com
 AEZ 2021; 2(1): 61-72
 Received: 02-11-2020
 Accepted: 28-12-2020

Hanan Salah El-Din Taha
 Central Agricultural Pesticide
 Laboratory, Dokki, Giza,
 Egypt

Survivorship assessment for *Spodoptera littoralis* susceptible and field population after insecticide sublethal dose treatment

Hanan Salah El-Din Taha

DOI: <https://doi.org/10.33545/27080013.2021.v2.i1a.32>

Abstract

Spodoptera littoralis (Lepidoptera noctuidae) (Boisd) is the most economic and series pest in Egypt with highly fecundity and dangerous attacking many crops. Insecticide efficiency trials showed failure and need more investigation periodically by many subject area. Susceptible laboratory reared population and field collected batches were subjected to bioassay of time mortality of the life survivors recording is the data producer by treating larvae with sublethal dose (LC₃₀) of insecticides candidates for this testing. Time-mortality data are improved method involves proportional survival and hazards modeling, known as Cox regression. The survival analysis of *S. littoralis* treated with sublethal dose of some insecticides exhibit slight differences in the median survival time (LT₅₀) observed for each insecticide, where the higher values was 224.4 hour for metaflumezone and the smallest was 170.1 for pyridalyl in susceptible population while in field, the higher was 336 for indoxacarb and the smallest was 144 hour for imidacloprid. The median survival time was 265 for susceptible control and 375 hour for field control. Log rank test exhibit significant differences among treatments for susceptible (Log-rank test, $Z = 259.91$, d.f. = 4, $p < 0.001$) and field (Log-rank test, $Z = 297.48$, d.f. = 4, $p < 0.001$). Hypothesis testing were computed for testing the survival curves, and every two hazard rates equality, at all times less than the maximum observed. Survival curves within insecticide groups and within population and control were differ according to p-values are less than 0.05, verify the null hypothesis is success to be rejected. Some hazard rates decreases with time in both population. The hazard ratios of insecticides in relation to control was 95% confidence intervals of the susceptible and field calculated were smaller than one prove great indication that field population were dying at a significantly slower rate than susceptible, and males population of some insecticides have more chance to die at any point in time more than males from the susceptible. The results for field population were achieved only 1% of population survived at 555 hour, from 20 to 30 % at 336 hour, from 30 to 40% and from 40 to 50 % after treatment for all pesticide tested, where population stages were old moths, emerged moth, pupae and 5th instar larvae respectively. This result evidenced that all insecticide tested at low concentrations treatment through a long period of exposure, must stimulate a continuous mortality and the time mortality assay using Kaplan–Meier and Cox regression analysis methodology were accurate and provide clear decision about pesticide toxicity, efficiency status and differences.

Keywords: *Spodoptera littoralis*, sublethal dose, economic

Introduction

The continually pest exposure to a variety of insecticides were associated with rapid development of insecticide resistance because of selection pressure intensity in agricultural eco-systems. IRM are more flexible department methods for insecticide operations, besides a new active ingredients introduction or increasing insecticide diversity in space and time. The long-lasting insecticide products considered a difficult tool in insecticide resistance management strategies options and application precisely in rotations and need more studies. The insecticide that proved to be insufficient control must be removed and switch to a different insecticide. The insecticide effect and strength on pest control assessment attains by different biological assays for measuring the direct the dose response relationship Sternberg and Thomas (2018) [28]. The need for analysing time to death data recently arises in a number of recommended pesticide used to control pests to compare the post treatment duration of the pest individual can still alive as effect of the lower insecticide dose efficiency to judge manipulation in charity. The Dose or concentration-mortality assay can determine the median lethal dose or concentration causes 50% of insect deaths at time-mortality experiments at a single point of insect deaths event recording after treatment during short

Corresponding Author:
Hanan Salah El-Din Taha
 Central Agricultural Pesticide
 Laboratory, Dokki, Giza,
 Egypt

time with varying amount of insecticide. Additional ways, is the repeated mortality measurements that count a time requirements of experiment (once-time- dose- mortality) give the lethal time LT_{50} which the time to kill 50 % of the given population. Throne *et al.*, (1995) ^[29], Preisler, and Robertson, (1992) take this kind of analysis as decisions about using logistic regression to analyse relationship between dose, mortality and between time to estimate LD_{50} and LT_{50} respectively. However, using cumulative mortality and the cumulative probability of survives according to administrating the predetermined definite dose of insecticide concentration treatment, after serial times of effects were more appropriate. Accurately until the end of the insect pest lifecycle, will give the median survival time from larvae treatment day until moths emerging and die to determine the sub lethal time (day) that kill 0.5 of the population. It can perform by using Aalen-Nelson Estimator) or the closely related Kaplan–Meier survival analysis estimates (Kaplan and Meier, 1958) ^[18]. This by assuming that: at any time pest individuals have the same survival predictions as others followed, the survival probabilities showing the mortality variation over time are the same for any insecticide and the event happens is the only varied at that time specified. Generally, it considered as specific methodology analysis for the detection of low-level insecticide resistance or insecticide poor efficiency Holbrook *et al.* 2002 ^[8]. Survival time are data can measure the time to a certain event such as death, failure, response, separation or the development of a given disease Brevik, *et al.*, (2018) ^[3] and Etikan *et al.*, (2017) ^[13]. Recently many papers stated that survival analysis is habitually uses in human public health studies to estimate how long a patient will survive fronting a disease and effects or compute the time to death following any hurt disease.

In this study, a Kaplan–Meier survival analysis accomplished to organize comparison between insecticides varied in their mode of action against two strains of *Spodoptera littoralis* (Boisd.) susceptible and field strain treated with candidate insecticide sublethal dose. To determine if insecticides by that dose can kill considerable number of a proper pest population attacking such crop, and how much number can pass to the next generation.

Materials and Methods

Insects and insecticides

Susceptible strain of cotton leaf worm *Spodoptera littoralis* (Boisd.) (Lepidoptera: noctuidae) used in this study mass reared in standard lab conditioned about many years in agriculture research center rearing laboratory rooms apart from insecticide exposure according to (Shaaban *et al.*, 1985) ^[27]. Otherwise Kalubia governorate field strain larvae of cotton leaf worm collected from cotton plant and used directly in bioassays alongside with the susceptible by applicant insecticides.

Insecticides used were provided by central agricultural pesticide laboratory as formulated materials. The chemical name, trade name, active ingredient percentage and field rate of the formulated insecticides were as follows: organophosphate compounds were chlorpyrifos (Tanner 48% EC, cm/l), dimethoate (Ares 40% EC, cm/l). Nicotinoid class was imidacloprid (Condorfast 35% SC, cm/l). Pyrethroid was cypermethrin (Biomethrin 10% EC, cm/l). Insect Growth Regulator was hexaflumeron (Dimeuron 10% EC, cm/l), lufenuron (Match 5% EC, cm/l).

Metaflumezone (Technomizone 22% SC, cm/l), novaluron (Roxy 10% EC, cm/l). Unclassified group of insecticide was pyridalyl (Peleo 50% EC, cm/l) and carbamate was Indoxacarb (Pret 15% EC, cm/l).

Concentration-mortality bioassays

Bioassays performed on all strains using a leaf dip method according to Paramasivam and Selvi (2017) ^[26] as proceeding. Uninfected cotton Leaves individually and efficiently dipped in subsequent diluted solutions of five insecticide concentration, and left to dry and controls dipped in water only. Afterward, treated leaves placed individually in 9-diameter petri dish filled with 10 individual of 1-day-old, 4th instars larvae and 3 replicate done for each concentration. The dishes preserved in controlled conditions at $25 \pm 2^{\circ}\text{C}$ and 16:8 h light: dark. Mortality was recorded 24 h or 48 h after treatment according the insecticide. Recorded leaf-dip bioassay data submitted to Polo Computer Software program for $LC_{30, 50, 90}$, estimation according to Finney (1971) ^[14], including correction of mortality.

Sublethal dose treatment

Sublethal effect of the insecticides on *S.littoralis* survivorship were performed by dipping cotton leaves in each insecticide solution containing the LC_{30} concentration, which previously determined for each insecticide by the baseline assay. After drying of the plant, leaves introduced to feed about 300 fourth instars larvae for each insecticide in one big container for 24 or 48 hour after that survived larvae collected and placed in new container and fed on fresh untreated leaves. Survived larvae recorded daily at constant time intervals until moth emergences, oviposition and die, then considered the end of the counts.

Survivorship assessment procedure

The mean idea of this test according to Goel *et al.*, 2010 ^[15] and Clark *et al.*, 2003 ^[6] is calculating the survival function (time) which is the probability that an individual will survive or die after a given time. The survival probability and the successive probabilities from one intervals to the next that can multiplying together to give cumulative survival probability, then the estimated probability refers to changes in value of each event in its group. The survival probability at each interval is calculated as the number of survivors divided by the number of individuals at risk by this formula: $S_t = \text{no. of individuals living at the start (n)} - \text{no. of individuals died(d)} / \text{no. of individuals living at the start}$. This step must followed by the measurement of median survival time (LT_{50}) according to Brookmeyer and Crowley (1982) ^[4] that means, the total probability of survival at the end of a time is 0.50. Alternatively, plotted directly from the survival plot lines. Survival function, according to Lee and Wang (2003) ^[20], is the probability that an individual will survive longer than a specified time t . The output of this analysis were included: firstly, the hazard ratio of the risk means the risk ratio that occurring at any time between groups. The ratio between $s(t)$ and $h(t)$ which is calculated by this formula $h(t) = -d/dt (\log s(t))$, and the cumulative hazard is $H_t = -\ln(S_t)$. The second output result were the log-rank test: is a statistical method to compare survival function and to test statistically difference effect of the survival times between two groups to death variable. Its calculation method as follows:

$$\chi^2(\log \text{rank}) = \frac{(O_1 - E_1)^2}{E_1} + \frac{(O_2 - E_2)^2}{E_2}$$

Where the O_1 and O_2 are the total no. of observed events between two group respectively, and E_1 and E_2 is the total no. of expected events. The expected no. of events calculated as the death risk at that time multiplied by the no. alive in the group. Under the null hypothesis, the death risk = (no. of deaths/no. alive). Finally the Cox proportion hazard test: is to test the probability of dying at given time as result of other independent variables effect, (Mantel and Haenszel, 1959; Cox, 1972) [21, 7] and test Confidence intervals calculated to prove that there is no significant difference between two groups. If test statistic value is less than the critical value (using chi square table) for degree of freedom.'

Statistical analysis

The data for serial time-death recording after predetermined sub lethal dose treatment (LC_{30} ppm) submitted to survival analysis statistic procedures estimator over the Kaplan-Meier (1958) [18] analysis time to event data for comparing group insecticides with control. All the analysis steps output was the non-parametric procedure (Survival or failure (the death event) probabilities and its confidence intervals, Log-rank test method, Hazard ratios and Cox regressions) at a definite times, and in addition to survival reliability curves, hazard rate and hazard ratio chart were performed using NCCSS2020 Computer Software (v.2019). The variable censor was coded zero if the event occurred at the exact time of the event occurrence. The control was omitted in the data set because there was no mortality for larva exposed to

water only.

Results and Discussions

Dose-mortality assay

The results of the *S.littoralis* susceptible population treated with insecticide tested and calculation was completed using the probit of Dose-mortality assays according to Finney probit analysis, that frequently used to detect the shift of response at one time of death recording are found in (Table 1). Generally LC_{50} values in ppm showed slight variations between insecticide classes and group and there was no death in control, which treated of water. In addition, kalubia field, is a different set of this pest population, result of LC_{50} and LC_{90} values, are found in (table 2). Toxicity lines fit showed low χ^2 values ranged from (0.16 to 4.3) and high p-values ranged from (0.11 to 0.96), proving accuracy to estimate the mortality curves for susceptible population. Indoxacarb had the highest toxicity followed by hexaflumuron, lufenuron, chlorpyrifos, pyridalyl, deltamethrin, novaluron, metaflumezone and imidacloprid with relative toxicity of 32.20, 739.43 and 1074.07 ppm, in susceptible population, respectively (Table 1). However, LC_{50} s of the field were the highest values was 329 ppm for imidacloprid and 58.2 ppm for dimethoate and the lowest was 3.7 ppm for hexaflumuron. Some insecticide had LC_{90} were lower than the field label rate and considered more effective. Statistical analysis of LC_{50} s significance proved no significant differences between insecticide each other and time but between susceptible population and time ($F=1.76$, $df=2$, $p=0.193$) and field between population and time ($F=2.3$, $df=2$, $p=0.055$). There are no significant differences between LC_{50} of field and susceptible strain.

Table 1: Probit analysis of *S. littoralis* susceptible population insecticide toxicity.

Insecticide	Slope± SE	LC_{30} (95% CI)	LC_{50} (95% CI)	LC_{90} (95% CI)	χ^2	pvalue	RT
Chlorpyrifos	1.76±0.17	13.86(10.4-17.19)	27.48(22.84-32.4)	146.3(111.3-214.3)	4.03	0.25	3.123
Dimethoate	1.58±0.20	29.1(21.3-36.3)	62.34(51.6-76.1)	400.7(262.2-794.3)	4.06	0.13	7.084
Imidacloprid	1.29±0.16	42.7(32.9-51.1)	108(92.4-132.7)	1051(609-2596.4)	1.78	0.409	12.273
Cypermethrin	1.8±0.11	3.3(2.0-5.4)	6.45(3.96-10.5)	32.9(20.2-53.6)	0.33	0.18	1.46
Hexaflumuron	1.86±0.21	4.6 (3.27-5.88)	8.86 (7.1-10.5)	43.1 (32.7-64.98)	4.3	0.11	1.007
Lufenuron	2.2±0.21	8.9(6.7-10.9)	15.4(12.7-17.9)	58.5(48.3-75.9)	3.9	0.14	1.750
Metaflumezone	1.78±0.15	31.3(25.7-36.9)	61.6(52.6-73.1)	322.6(235.4-497.7)	0.27	0.96	7.000
Novaluron	2.5±0.23	30.0(25.0-34.8)	48.2(42.2-54.8)	153.4(125.4-201.4)	3.5	0.16	5.477
Pyridalyl	1.68±0.17	21.3(16.5-26.0)	43.6(36.2-52.9)	251.6(177.1-415.9)	4.0	0.14	4.955
Indoxacarb	1.89±0.22	6.7(4.7-8.5)	12.6(10.2-15.0)	59.9(66.6-154.4)	0.16	0.92	1.432

SE= standard error of the slope from the logistic method. CI= confident intervals. Resistance ratios. RT= relative toxicity means the division of each LC_{50} by the less value of the susceptible population.

However, in field population p-values of the probit analysis were ranged from 0.027 to 0.78 for and χ^2 ranged from 1.07 to 6.1. The resistance ratios were represent dissimilar values (ranged from 0.448 to 1.429) and differed from each other (Table 2). The highest RR were 7 for imidacloprid then 1.429 was for indoxacarb and the lowest was 0.42 for pyridalyl. When comparing LC_{50} values with field rate some

insecticide LC_{50} like indoxacarb, lufenuron and chlorpyrifos slightly are not enough to kill 50% of field population (Table 2). Nevertheless, other insecticide were below the field rate recommended concentration (insecticide lable), means this strain were very susceptible to those insecticide that considered newest chemical products used to control crop pests.

Table 2: Probit analysis of *S. littoralis* field population larvae insecticide toxicity.

Insecticide	Slope± SE	LC_{30} (95% CI)	LC_{50} (95% CI)	LC_{90} (95% CI)	χ^2	Pvalue	RR
Chlorpyrifos	1.14±0.15	11.3 (5.4-17.9)	32.7 (21.5-43.6)	433.0 (287.9-827.1)	1.07	0.78	1.190
Dimethoate	1.86±0.16	30.4 (24.0-36.74)	58.2 (49.3-67.8)	283.6 (221.3-395.1)	3.4	0.33	0.934
Imidacloprid	2.5±0.347	177.6(125.2-226.7)	329.9(262.4-411.9)	1069.9(783.8-1748.6)	1.22	0.74	7.0
Cypermethrin	1.94±0.11	13.9(8.0-21.2)	25.1(15.4-40.8)	124.0(76.2-201.1)	0.45	0.71	1.54

Hexaflumeron	0.98±0.16	1.16 (0.31-2.3)	3.97 (1.888-6.0)	80.1 (48.7-200.4)	3.53	0.31	0.448
Lufenuron	2.55±0.25	9.59 (7.5-11.48)	15.4 (13.0-17.6)	48.8 (40.7-62.7)	4.5	0.104	1.000
Metaflumezone	1.57±0.187	18.9 (12.2-20.8)	33.8 (25.6-41.7)	221.0 (164.1-344.8)	1.7	0.89	0.549
Novaluron	1.8±0.24	13.13 (7.9-18.0)	25.6 (18.8-31.7)	131.75 (100.9-200.4)	4.0	0.135	0.531
Pyridalyl	0.95±0.10	5.14 (2.47-8.5)	18.3 (11.7-25.5)	404.6 (269.5-718.6)	2.86	0.412	0.420
Indoxacarb	2.6±0.20	11.37 (6.4-15.0)	18.0 (11.8-25.4)	55.45 (42.9-112.1)	6.1	0.027	1.429

SE= standard error of the slope from the logistic method. CI= confident intervals. Resistance ratios (RR) = LC_{50S} and LC_{95S} of the four field strains, divided by the LC_{50} and LC_{95} of the lab strain

Time mortality assay

The survival and failure probabilities and the proportional hazards (Cox) modelling to analyze time-to-event data were calculated to obtain different result about determining the percentage of mortality through series of time intervals records of larvae death and the rate of death alongside the pest lifecycle this results were in Table (3 and 4). This test including the cumulative survival probabilities, cumulative hazard function, and hazard rates, all of those represent the

survival analysis machineries and the life table that constructed from a set of grouped of mortality failure and censor data. In addition to the hazard, ratios that used to compare directly the death rate of one group to that of a control group, similar to resistance ratios that used to assess insecticide resistance in the field populations that exhibit heterogeneity and such resistance phenotypes, these results were in (Table 5, 6, 7 and 8).

Table 3: Estimated Survival and its confidence Limits report for some insecticide sublethal toxicity against *S.littoralis* susceptible and field population.

Insecticide	Susceptible population			Field population		
	Larvae	Pupae	Moth	Larvae	Pupae	Moth
Chlorpyrifos	0.43(0.40-0.46)	0.24(0.22-0.27)	0.026(0.016-0.035)	0.27(0.24-0.30)	0.135(0.11-0.16)	0.13(0.005-0.002)
Dimethoate	0.44(0.41-0.47)	0.23(0.21-0.26)	0.02(0.013-0.03)	0.27(0.24-0.31)	0.14(0.11-0.16)	0.014(0.005-0.02)
Imidacloprid	0.469(0.44-0.5)	0.27(0.24-0.29)	0.03(0.02-0.039)	0.25(0.22-0.28)	0.13(0.10-0.153)	0.015(0.002-0.018)
Cypermethrin	0.43(0.39-0.46)	0.24(0.21-0.27)	0.025(0.016-0.036)	0.36(0.33-0.4)	0.2(0.17-0.22)	0.016(0.008-0.025)
Hexaflumero	0.46(0.43-0.49)	0.24(0.22-0.26)	0.018(0.011-0.027)	0.34(0.31-0.38)	0.19(0.16-0.21)	0.019(0.0097-0.0029)
Metaflumezone	0.51(0.49-0.54)	0.31(0.28-0.34)	0.033(0.023-0.043)	0.35(0.31-0.38)	0.16(0.13-0.18)	0.012(0.004-0.02)
Lufenuron	0.47(0.44-0.56)	0.25(0.22-0.28)	0.019(0.012-0.028)	0.33(0.3-0.37)	0.18(0.15-0.2)	0.02(0.008-0.03)
Indoxacarb	0.47(0.44-0.5)	0.26(0.24-0.29)	0.025(0.016-0.035)	0.36(0.33-0.4)	0.17(0.15-0.2)	0.017(0.008-0.025)
Thiametham	0.47(0.44-0.496)	0.28(0.25-0.304)	0.03(0.021-0.041)	0.33(0.3-0.36)	0.17(0.16-0.2)	0.019(0.009-0.027)
Pyridalyl	0.39(0.354-0.42)	0.17(0.14-0.193)	0.014(0.006-0.02)	0.32(0.28-0.36)	0.17(0.14-0.2)	0.017(0.007-0.27)
Control	0.61(0.59-0.63)	0.39(0.37-0.41)	0.04(0.04-0.052)	0.6(0.58-0.62)	0.39(0.36-0.4)	0.04(0.034-0.05)

Censored is the missing individual and failure is the individual failed to die and survived to the next time intervals.

The overall dead or individual didn't survive LC_{30} treatment after experiment finished was 74,72,70,68,78,74,76,56,73 and 72 % and 90,90,93,85,87,85, 86,80,85, and 88 % in susceptible and field population respectively, for chlorpyrifos, dimethoate, imidacloprid, thiamethoxam, lufenuron, hexaflumeron, cypermethrin, metaflumezone, indoxacarb, and pyridalyl respectively, as shown in commulative individual number at risk inside analysis. The mortality rate and the Kaplan-Meier survival curves of *S.littoralis* larvae treated with the different insecticide classes were not significantly different ($P=0.00$) from the control. The analysis of mortality rates also revealed that the effects of insecticide treatments within insect stages were ($F = 1.189$, $df = 3$, $P = 0.306$) in the susceptible population, but in the field population ($F = 0.642$, $df = 2$, $P = 0.427$) and the interaction between pest stages (Larvae-Pupae and Moth) were highly significant for the ($F = 1.189$, $df = 3$, $P = 0.306$).

Generally, Survival was significant and affected by insecticide dose. The order and death rate in the two population was not always the same trend (Fig. 1, 2). Compare the daily commutative survival pattern for insecticide treatment groups had different probability of surviving with a control, while another took longer time to deaths. The survival analysis showed that the order and patterns of the lines of cumulative survival for the various insecticides were significant in both assays, Kurtosis Normality was -0.363 for susceptible but for field was -0.813 and p-values for susceptible was 0.716 and for field was 0.415. Susceptible population showed survival curve lines were very closer or overlying also showed statistically no significant differences and no clear pattern ($p=0.0$) in the rates (Skewness Normality 4.59). On the dissimilarity, field population showed divergent survival line (Skewness Normality 5.578) (Table 7). The dynamics of survival, observed increase of mortality caused an increase of the death rate because of serious damage to treated larvae through experiment duration (Fig 3, 4).

Table 4: Estimated Hazard and its confidence Limits report for some insecticide sublethal toxicity against *S.littoralis* susceptible and field population.

insecticide	Susceptible population			Field population		
	Larvae	Pupae	Moth	Larvae	Pupae	Moth
Chlorpyrifos	0.84(0.77-0.92)	1.41(1.30-1.52)	3.68(3.33-4.10)	0.006(0.0054-0.0069)	0.004(0.003-0.005)	0.0001(0.00150-0.0017)
Dimethoate	0.82(0.75-0.89)	1.32(1.22-1.42)	3.89(3.51-4.35)	0.0057(0.005-0.0065)	0.003(0.0024-0.0039)	0.0014(0.0012-0.0016)
Imidacloprid	0.76(0.7-0.82)	1.3(1.22-1.41)	3.55(3.24-3.92)	0.0051(0.0044-0.0059)	0.0016(0.001-0.0023)	0.0012(0.01-0.0014)
Cypermethrin	0.85(0.78-0.93)	1.42(1.32-1.54)	3.70(3.35-4.15)	0.0035(0.0031-0.0041)	0.001(0.0008-0.0016)	0.0015(0.0013-0.0017)
Hexaflumeron	0.78(0.72-0.85)	1.431.33-1.54)	4.0(3.61-4.492)	0.0047(0.0003-0.0041)	0.0004(0.003-0.0045)	0.0015(0.0013-0.0017)

Metaflumezoxon	0.67(0.62-0.73)	1.18(1.1-1.26)	3.42(3.13-3.75)	0.0035(0.003-0.004)	0.0032(0.0025-0.004)	0.0014(0.0012-0.0016)
Lufenuron	0.76(0.69-0.82)	1.4(1.32-1.54)	3.95(3.57-4.4)	0.0043(0.0037-0.005)	0.0028(0.0022-0.0035)	0.0014(0.0012-0.0016)
Indoxacarb	0.76(0.69-0.82)	1.34(1.24-1.44)	3.7(3.36-4.12)	0.002(0.0017-0.0024)	0.0035(0.0029-0.0043)	0.001(0.0001-0.0013)
Thiametham	0.77(0.70-0.83)	1.29(1.19-1.39)	3.51(3.19-3.89)	0.0052(0.0046-0.0058)	0.004(0.003-0.0048)	0.0013(0.0012-0.0015)
Pyridalyl	0.95(0.87-1.04)	1.79(1.65-1.9)	4.47(3.93-5.21)	0.0043(0.0037-0.005)	0.0039(0.0032-0.0047)	0.0012(0.0012-0.0013)
Control	0.5(0.47-0.54)	0.95(0.89-0.99)	3.14(2.96-3.35)	0.0001(0.0001-0.005)	0.0042(0.0034-0.0051)	0.0002(0.0001-0.0001)

Censored is the missing individual and failure is the individual failed to die and survived to the next time intervals.

Some hazard rates decreases with time, others have a hazard rate that increases with time, some are constant, and some exhibit all three behaviors at different points in time. In fact, the subjects who die (fail) as well as subjects who are censored or not presented, the program survival datasets and computation required the start and the end of survival times and observations of censored or failed to perform several logrank tests and provides both the non-parametric and randomization test significance levels. The dataset contained 24-observation time (number of events) for each insecticide groups passing from the larvae, pupae, and moths that

censored or failed. The results for field population were achieved only 1% of population survived at 555 hour, from 20 to 30 % at 336 hour, from 30 to 40% and from 40 to 50 % after treatment for all pesticide tested, where population stages were old moths, emerged moth, pupae and 5th instar larvae respectively. With observation of recording individuals escaping or dying accidentally, the delayed toxicity of insecticide tested may be reason of the differential sensitivity of *S. littoralis* according to age distribution and the accumulation of the insecticide compounds in bodies' leads to sensitivity variation. This result evidenced that all insecticide tested at low concentrations treatment through a long period of exposure, must stimulate a continuous mortality.




Fig 1: Susceptible and field population of *S. littoralis* status of surviving insecticide treatment, box plot differentiation between insecticides and population through comparing the center and spread of groups. The spread look differences and fill weight of ten insecticide treatment boxes. Median weight of groups not similar and the general weight of them more variable than other. Also there is differences between the centers of the group. The number above were Identifying the cause of outlier. Skewed data, the majority of data located in high side of the graph. Skewness indicated that data may be not normally distributed and refer to failure time, and some data Fail to die closely and more fail later.

A slight differences in the median survival time (LT₅₀) observed for each insecticide, where the higher values was 224.4 hour for metaflumezone and the smallest was 170.1 for pyridalyl in susceptible population while in field, the higher was 336 for indoxacarb and the smallest was 144 for imidacloprid. The median survival time was 265 for susceptible control and 365.9 hour for field control (Table 5). The survival analysis of *S. littoralis* treated with sublethal dose of some insecticides indicated (Table 3, 4) exhibit significant differences among treatments for

susceptible (Log-rank test, Z = 259.91, d.f. = 4, p < 0.001) and field (Log-rank test, Z = 297.48, df. = 4, p < 0.001) (Table 6). The survival of larvae was 100% in the control (without insecticide exposure) after 500 h of exposure. This result were similar to De Castro *et al.*, 2013 [9], calculate high median survival time (LT₅₀) of predatory stinkbugs *Podisus nigrispinus* (hemipteran: pentatomidae) exposed chlorantraniliprole, methamidophos, spinosad, deltamethrin. However, when exposed to dried insecticide residues indicated significant differences among treatments.

Table 5: Median survival time and Restricted Mean Survival Time of the two population from Nonparametric Kaplan-Meier test.

Insecticides	Susceptible population				Field population			
	Median (CL95%)	RMST (τ) (CL95%)	±SE	MXL.	Median (CL95%)	RMST(τ)	±SE	MXL.
Chlorpyrifos	193.983(192-216)	251.4(241.3-261.6)	5.159	191.87	168(144-144)	207.9(195.9-220)	6.14	164
Cypermethrin	191.158(192-216)	251.7(241.1-262.3)	5.390	189.18	192(168-192)	258.5(245.3-271.8)	6.75	254
Dimethoate	195.325(216-216)	245.1(236.0-254.2)	4.639	192.33	168(144-144)	214.4(201.3-227.5)	6.67	163
Metaflumezone	224.407(240-240)	280.2(270.6-289.7)	4.890	218.28	240(216-216)	271.7(257.9-285.6)	7.07	237
Hexaflumeron	200.702(216-216)	248.6(239.8-257.4)	4.496	196.48	192(168-168)	235.4(222.6-248.2)	6.54	190
Imidacloprid	208.825(240-240)	279.1(269.0-289.1)	5.152	204.89	144(144-144)	219.8(205.1-234.5)	7.48	141
Indoxacarb	204.912(216-216)	266.5(256.5-276.4)	5.064	200.74	336(288-288)	313.3(298.5-328.1)	7.56	335
Lufenuron	204.924(216-216)	251.6(243.0-260.2)	4.393	200.27	168(144-168)	230.8(216.6-245)	7.23	163
Thiamethoxam	206.313(216-216)	266.6(256.2-277.0)	5.316	202.49	168(168-168)	228.2(216.2-240.2)	6.12	163
Pyridalyl	170.107(192-192)	222.2(212.7-231.6)	4.831	168.35	168(144-168)	230.2(216-244.4)	7.24	162
Control	265.946	508.7(503-514.3)	2.890	257.88	356.9	508.7(503-514.3)	2.89	


Median Estimates are based on an interval upper limit of τ = 552.0, the maximum observed time among all groups.

MXL=Nonparametric test or Maximumlikelihood estimated for mean and median survival time.

RMST (t)= Restricted Mean Survival Time. ±SE= Standard error of the mean.

The logrank test is testing the equality of hazard functions or estimated survival times. This test uses W (T) = one, that is, equal weighting, optimum power when the hazard rates

of the K populations are proportional to each other which is only valid for equal censoring (Lee and Wang (2003) [20]).






Fig 2: Susceptible and field survival curves from Kaplan Meier estimates determined the serial interval time to death alongside experiments duration from the day 1 of the 4th instar larvae to moth emergence, oviposition and die. Censoring and the survival probability, number at risk, included for plotting curves, median survival time can be pointed out. Survival data are non-normal (skewed) so no distributional assumption. Survivorship curve result here of *S.littoralis* lifecycle, all are fall between one and two type, Pearl (1941) [25].

Survival of any pest population normally depends on, exposure; pressure of multiple stressors like agro-chemicals, pathogens and poor food resources those causing population reductions. Bunya *et al.*, 2016 [5], there was a decrease in survival at a parathion dose 20 mg/kg, equal one-half of rat animals survived to 24 hours and to 8 days after poisoning, prove dose dependent calculated survival rate was 75, 79%

per day of sixth instars fall armyworm treated with azinophosmethyl and alphacypermethrin (Ali and Lutterell 1990) [2]. Matos, *et al.*, 2017 [22], in bed bugs (*Cimex lectularius* L.) fed on fipronil or clothianidin,, Nymph survivorship was depend on the presence of fed or unfed adults.

Table 6: Log-rank statistic and hazard rate for insecticide time-mortality assay of the *S.littoralis* susceptible and Field populations.

Insecticides	Susceptible population			Field population		
	Z-Value	Standard Error	Standardized Z-Value	Z-Value	Standard Error	Standardized Z-Value
Chlorpyrifos	168.075	23.623	7.115	229.282	16.553	13.851
Cypermethrin	160.985	22.913	7.026	117.08	19.087	6.134
Dimethoate	255.322	23.906	10.680	203.096	16.339	12.43
Metaflumezone	58.300	27.171	2.146	60.539	18.553	3.263
Hexaflumeron	293.816	24.350	12.066	175.985	17.927	9.817
Imidacloprid	34.060	25.548	1.333	171.032	15.564	10.989
Indoxacarb	126.666	24.801	5.107	-49.332	19.333	-2.552
Lufenuron	296.929	25.049	11.854	158.262	16.423	9.636
Thiamethoxam	82.005	24.784	3.309	207.743	18.277	11.366
Pyridalyl	277.670	20.173	13.764	156.576	16.413	9.54
Control	-1753.828	37.320	-46.994	-1430.265	31.948	-44.768

The logrank test is testing the equality of hazard functions or estimated survival times. The test imply for susceptible, Chi-Square = 2257.250, DF = 10, Probability Level = 0.0000, Randomization Test Probability Level = 0.0050 (200 Monte Carlo Samples) Weighting of Hazard Comparisons across Time = Equal. For field, Chi-Square = 2414.278, DF = 10, Probability Level = 0.0000 Randomization Test Probability Level = 0.0050 (200 Monte Carlo Samples) Weighting of Hazard Comparisons across Time = Equal.

For testing the survival curves, and if the hazard rates, of two or more populations are equal, at all times less than the maximum observed time, hypothesis testing were computed. The alternative hypothesis is that at least two of the hazard


rates are different at some time less than the observed maximum time (Table 7). The null hypothesis means that there is no difference between the groups of insecticide treatment.

Table 7: One-Sample T-Test Report, Tests of Assumptions.

Different Test of Assumption	Susc. population			Field population		
	Value	Prob Level	Decision ($\alpha = 0.050$)	Value	Prob Level	Decision ($\alpha = 0.050$)
Shapiro-Wilk Normality	0.913	0	Reject normality	0.836	0	Reject normality
Skewness Normality	4.591	0.000004	Reject normality	5.578	0	Reject normality
Kurtosis Normality	-0.363	0.716326	Cannot reject normality	-0.813	0.415921	Cannot reject normality
Omnibus Normality	21.213	0.000025	Reject normality	31.784	0	Reject normality
Martinez-Iglewicz	1.039	0	Reject normality	0.955	0	Can't reject normality
Kolmogorov-Smirnov	0.119	0	Reject normality	0.097	0	Reject normality
D'Agostino Skewness	-33.184	0	Reject normality	-20.50	0	Reject normality
D'Agostino Kurtosis	-7.96	0	Reject normality	-25.11	0	Reject normality

Table (7), shows that p-values are less than 0.05, this means that the null hypothesis is success to be rejected. Then the survival curves of within insecticide groups and within population and control were differ. Holbrook, *et al.*, 2002^[8], the detection of low levels of fipronile resistance in the

German cockroach using logistic regression and resistance ratios are relatively ineffective in distinguishing differences between strains, but dose-mortality or time-mortality data are improved method involves proportional hazards modeling, known as Cox regression.






Fig 3: The probability distribution of individuals at time T of certain events itself could roll the event of the next prompt generating conditional failure rate in reliability this is the Hazard rate, $h(T)$: or the age specific failure rate, that has a different hazard rate function. It is essential in computing and quantifying events in survival analysis.

Table 8: Hazard ratio-Cox-Mantel Logrank Test in relation to control (Cox-Mantel Logrank Test).

Group Ratio	Susc. population					Field population				
	Hazard Ratio	Chi-Square	Prob Level	Lower	Upper	Hazard Ratio	Chi-Square	Prob Level	Lower	Upper
control/Chlorpyrifos	0.08	1627.194	0.0000	0.07	0.09	0.07	1786.251	0.0000	0.06	0.09
control/Cypermethrin	0.08	1615.426	0.0000	0.07	0.09	0.09	1250.187	0.0000	0.08	0.11
control/Dimethoate	0.07	1884.783	0.0000	0.06	0.08	0.07	1656.557	0.0000	0.06	0.09
control/Metaflumezone	0.10	1406.291	0.0000	0.08	0.11	0.1	1092.277	0.0000	0.08	0.12
control/Hexaflumeron	0.07	2023.117	0.0000	0.06	0.08	0.08	1505.797	0.0000	0.07	0.1
control/Imidacloprid	0.10	1320.451	0.0000	0.09	0.11	0.08	1488.066	0.0000	0.06	0.1
control/Indoxacarb	0.09	1539.402	0.0000	0.08	0.10	0.13	751.388	0.0000	0.11	0.16
control/Lufenuron	0.07	2011.115	0.0000	0.06	0.08	0.08	1465.173	0.0000	0.07	0.1
control/Thiamethoxam	0.09	1415.214	0.0000	0.08	0.11	0.08	1618.578	0.0000	0.07	0.09
control/pyridalyl	0.07	2030.946	0.0000	0.06	0.08	0.08	1454.114	0.0000	0.07	0.1

Hazard ratio is the relative survival between groups garnered from proportional hazards models and if less than one it refer to that individual males of a field population were dying more slowly than the susceptible population. The results concluded about time mortality assay using Kaplan–Meier and Cox regression analysis methodology were accurate and provide clear decision about pesticide toxicity elucidation, pesticide efficiency status and differences. The hazard ratios of insecticides in relation to control and their 95% confidence intervals of the susceptible and field calculated were smaller than one (Table 7). This is a great indication that field population were dying at a significantly slower rate than susceptible. Males of the

indoxacarb insecticide in field were found to be about one-third (hazard ratio: 0.13/0.09) as possible to die at any point in time more than males from the susceptible. Some theoretical demonstrations about survival argument after organism exposed to stress like insecticides ranged from low to high toxic effect, some of it as follows. The wood frog 15 populations exposed to carbaryl, and tolerance pattern noticed in consistency with theoretical predictions, that populations was closed to agriculture displayed constitutive tolerance to carbaryl whereas populations far from agriculture had low native tolerance but high magnitudes of induced tolerance (Hua, *et al.*, 2015).





Fig 4: The hazard function or death rate is the prompt probability of deaths for individuals that still alive.

Papachristos, *et al.*, 2008 ^[12], study the of the Mediterranean fruit fly, *Ceratitis capitata* (Wiedemann) (Diptera: Tephritidae), immature stages on the lemon, and bitter oranges had strong effects on larval performance and survival, essence of the peel was the most critical, inducing very high mortality rates. Survival of larvae ranged from zero for all varieties tested to 22.5%, except for bitter orange (76%). Survival of pupae was high for all varieties tested (81.1-90.7%). Kingsolver *et al.*, 2011 ^[17], study the changes in temperature-dependence of *Manduca sexta* population growth can affect juvenile survival and adult reproduction. High temperatures prevent survival to maturity. The differing in responses of multiple life stages is essential to understand the ecological and evolutionary

consequences of climate change. Lashkari, *et al.*, 2007 ^[19], found no significant differences in aphid life-table parameters between the two imidacloprid and pymetrozine treatment on population growth parameters of cabbage aphid, *Brevicoryne brassicae* L. (Homoptera: Aphididae). Cox regression coefficient, was computed to the time-mortality data using logistic models for all insecticide fit model (ranged from 0.0061 to 0.46 for susceptible and -0.17 to 0.47 for field), of the susceptible and field strains. If the coefficient was negative means the insecticide was very effective on the pest, but the maximum value means the insecticide loss efficiency to control the pest. The regression coefficient were supported in itself, with significant Wald statistic consists of Z test, probability level, pseudo R².

Table 9: Regression Coefficient (B) of Cox proportion hazard test model for *S.littoralis* susceptible and field population mortality analysis taking risk ratio, in relation to pyridalyl insecticide.

Insecticide	Susceptible Population						Field Population					
	Coefficient (B)±SE	Risk Ratio Exp(B)	Mean	Wald Z-Value	Prob Level	Pseudo R ²	Coefficient (B)±SE	Risk Ratio Exp(B)	Mean	Wald Z-Value	Prob Level	Pseudo R ²
Chlorpyrifos/	0.1408±0.039	1.151	0.012	3.59	0.0003	0.0014	0.439±0.044	1.55	0.004	9.96	0	0.0188
Cypermethrin/	0.006±0.045	1.006	0.007	0.13	0.8934	0	0.257±0.042	1.29	0.017	6.12	0	0.0072
Dimethoate/	0.319±0.034	1.37	0.015	9.43	0	0.0099	0.357±0.046	1.42	0.003	7.62	0	0.0111
Metaflumezon/	0.311±0.0317	1.36	0.026	9.80	0	0.0107	-0.178±0.046	0.83	0.015	-3.81	0.0001	0.0028
Hexaflumeron/	0.399±0.032	1.49	0.016	12.38	0	0.0169	0.38±0.0459	1.46	0.009	8.29	0	0.0131
Imidacloprid/	0.011±0.035	1.01	0.020	0.33	0.7407	0	0.191±0.052	1.21	-0.000	3.66	0.0002	0.0026
Indoxacarb/	0.111±0.035	1.11	0.017	3.13	0.0017	0.0011	-0.546±0.05	0.57	0.020	-10.75	0	0.0219
Lufenuron/	0.469±0.030	1.59	0.020	15.37	0	0.0258	0.307±0.057	1.36	-0.000	5.38	0	0.0056
Thiamethoxam/	0.098±0.038	1.10	0.015	2.55	0.0105	0.0007	0.475±0.04	1.60	0.013	11.71	0	0.0258
Control	-1.922±0.155	0.14	0.107	-12.36	0	0.0168	-1.96±0.173	0.139	0.162	-11.3	0	0.0242

Cox-Mantel and Mantel-Haenszel Logrank Tests; the Low Rank refer to events happening later in time.

Many literature cited such results about insect exposed to stress as honeybee workers exposed to sub-lethal doses of diet containing thiacloprid show higher survival rates was 11 days post exposure than controls Dickel *et al.*, (2018) ^[10]. Coulon *et al.*, (2018) ^[8], found statistical differences between survival curves according to Kaplan-Meier estimation between control bees and bees exposed to 2.5 ng/bee/day of thiamethoxam. Also combined neonicotinoid pesticide and parasite stress to honeybee queens' alter physiology and survival (Dussaubat *et al.*, 2016) ^[11]. All literature agree with the interpretation of the variation in survival observation within a population return to the biochemical process responsible for insecticide insect body eliminations or detoxifications.

References

- Aalen OO, Johansen S. An Empirical Transition Matrix for Non-Homogeneous Markov Chains Based on Censored Observations. *Scandinavian Journal of Statistics* 1978;5:141-150.
- Ali A, Lutterell RG. Survival of fall armyworm (Lepidoptera: noctuidae) immatures on cotton. *Florida entomologist* 1990;73(3):459-465.
- Brevik K, Schoville SD, Mota-Sanchez D, Chen YH. Pesticide durability and the evolution of resistance: A novel application of survival analysis. *Pest Manag Sci* 2018;74:1953-1963.
- Brookmeyer R, Crowley J. A confidence interval for the median survival time. *Biometrics* 1982;38:29-41.
- Bunya N, Sawamoto K, Benoit H, Bird SB. The Effect of Parathion on Red Blood Cell Acetylcholinesterase in the Wistar Rat. *Journal of Toxicology*, 2016, Article ID 4576952, 5 pages.
- Clark TG, Bradburn MJ, Love SB, Altman DG. Survival Analysis PartI: Basic concepts and first analyses. *British Journal of Cancer* 2003;89:232-238.
- Cox DR. Regression models and life tables. *Journal of the Royal Statistical Society* 1972;B34:187-220.
- Coulon MF, Schurr Martel AC, Cougoule N, Bégaud A, Mangoni P, Dalmon A *et al.* Survival analysis for the detection of low-level insecticide resistance. *Proceeding of the 4th international conference on urban pests 2002*, 159 -166.
- De Castro AA, Corrêa AS, Legaspi JC, Guedes RNC, Serrão JE, Zanuncio JC. Survival and behavior of the insecticide-exposed predators *Podisus nigrispinus* and *Supputius cincticeps* (Heteroptera: Pentatomidae). *Chemosphere* 2013;93:1043-1050.
- Dickel F, MuËnch D, Amdam GV, Mappes J, Freita D. Increased survival of honeybees in the laboratory after simultaneous exposure to low doses of pesticides and bacteria. *PLoS ONE* 2018;13(1):e0191256.
- Dussaubat C, Maisonnasse A, Crauser D, Tchamitchian S, Bonnet M, Cousin M *et al.* Combined neonicotinoid pesticide and parasite stress alter honeybee queens' physiology and survival. *Sci. Rep* 2016;6:31430.
- Dimitrios P, Papachristos Nikos T, Papadopoulos George Nanos D. Survival and Development of Immature Stages of the Mediterranean Fruit Fly (Diptera: Tephritidae) in Citrus Fruit. *J. Econ. Entomol* 2008;101(3):866-872.
- Etikan İ, Abubakar S, Alkassim R. The Kaplan Meier Estimate in Survival Analysis. *Biom Biostat Int J* 2017;5(2):00128.

14. Finney DJ. Probit Analysis. Cambridge University Press 1971.
15. Goel MK, Khanna P, Kishore J. Understanding survival analysis: Kaplan-Meier estimate. *Int J Ayurveda Res* 2010;1(4):274-278.
16. Hua J, Jones DK, Mattes BM, Cothran RD, Relyea RA, Hoverman JT. The contribution of phenotypic plasticity to the evolution of insecticide tolerance in amphibian populations. *Evolutionary Applications* published by John Wiley & Sons Ltd 2015;8:586-596.
17. Joel G, Kingsolver H, Woods A, Buckley LB, Potter KA, MacLean HJ *et al.* Complex Life Cycles and the Responses of Insects to Climate Change. *Integrative and Comparative Biology* 2011;51(5):719-732.
18. Kaplan EL, Meier P. Nonparametric estimation from incomplete observations. *Journal of the American Statistical Association* 1958;53:457-81.
19. Lashkari MR, Sahragard A, Ghadamyari M. Sublethal effects of imidacloprid and pymetrozine on population growth parameters of cabbage aphid, *Brevicoryne brassicae* on rapeseed, *Brassica napus* L. *The Authors Insect Science* 2007;14:207-212.
20. Lee ET, John Wang W. *Statistical Methods for Survival Data Analysis*, Third Edition. Book Series: Wiley Series in Probability and Statistics 2003.
21. Mantel N, Haenszel W. Statistical aspects of the analysis of data from retrospective studies. *Journal of the National Cancer Institute* 1959;22:719-748.
22. Matos YK, Sierras A, Schal C. Evaluation of the Potential for Secondary Kill for Ingested Insecticides in the Common Bed Bug (Hemiptera: Cimicidae). *Journal of Economic Entomology* 2017;110(3):1218-1225.
23. NCSST-2019, Statistical Software. NCSST, LLC. Kaysville, Utah, USA 2019. ncss.com/software/ncss.
24. Nelson W. A Short Life Test for Comparing a Sample with Previous Accelerated Test Results. *Technometrics* 1972;14:175-185.
25. Pearl A, Park T, Miner JR. Experimental studies on the duration of life. XVI. Life tables for the flour beetle 1941.
26. Paramasivam M, Selvi C. Laboratory bioassay methods to assess the insecticide toxicity against insect pests-A review. *Journal of Entomology and Zoology Studies* 2017;5(3):1441-1445.
27. Shaaban AM, Abo-Elghar MR, Abdel-Mohymen MR, El-Malla MA. Resistance of the Egyptian cotton leafworm, *Spodoptera littoralis* (Boisd.), to certain insecticides. *Zeitschrift für Pflanzenkrankheiten und Pflanzenschutz / Journal of Plant Diseases and Protection* 1985;92(1):69-75.
28. Sternberg ED, Thomas MB. Insights from agriculture for the management of insecticide resistance in disease vectors. *Evolutionary Applications* 2018;11:404-414.
29. Throne JE, Weaver DK, Chew V, James Baker E. Probit analysis of correlated data: multiple observations over time at one concentrate ion. *Journal of Economic Entomology* 1995;88(5):1510-1512.