

E-ISSN: 2708-0021  
 P-ISSN: 2708-0013  
[www.actajournal.com](http://www.actajournal.com)  
 AEZ 2022; 3(1): 15-18  
 Received: 07-11-2021  
 Accepted: 14-12-2021

**Mark Cooper**  
 School of Animal, Plant &  
 Environmental Sciences,  
 University of the Witwatersrand,  
 Johannesburg, South Africa

## Does sexual size dimorphism vary with female size in forest millipedes *Centrobolus* Cook, 1897?

**Mark Cooper**

DOI: <https://doi.org/10.33545/27080013.2022.v3.i1a.57>

### Abstract

This study aimed to determine whether the variation responsible for breaking Rensch's Rule when Sexual Size Dimorphism (SSD) increased with body size was due to male-biased or female-biased variation by studying a sex-based set of morphological characters. Variation in female volume based on length and width measurements was associated with the SSD in the forest millipede genus *Centrobolus*. There was a significant positive correlation between SSD and female size ( $r=0.56$ , Z score=2.76,  $n=22$ ,  $p<0.01$ ). The correlation coefficient between SSD and female size did not differ significantly from the correlation coefficient between SSD and species size (z-statistic=-0.7078,  $p=0.4790$ ). The morphological differences among the polygynandrous reproductive systems occurring in *Centrobolus* with higher SSD and larger females were discussed.

**Keywords:** Dimorphic, females, gradient, morphology, size, species.

### 1. Introduction

Numerous studies are finding animal taxa having female-biased SSD and mostly disobey Rensch's rule including corvids and pinnipeds [1, 2, 4, 13, 15, 18-19, 22-30, 34, 35, 37-40]. The finding of converse or inverse Rensch rule implies SSD increases with body size when females are larger [1, 2, 4, 13, 15, 18-19, 22-30, 34, 35, 37-40]. This has implications in the class Diplopoda because females are larger than males and SSD increases with body size [6-11].

The forest genus *Centrobolus* of the diplopod Order Spirobolida diplopod found along the east coast of southern Africa were the subject of this study. The millipede genus *Centrobolus* is located in the temperate region of South Africa, with its northern boundary on the east coast of southern Africa [15]. Millipedes of the *Centrobolus* genus break Rensch's rule and illustrate converse patterns of increasing female-biased sexual dimorphism variation with body size [6-11]. The problem with these and other tests of Rensch's rule is that they combine male and female variation among two sexes into species body size. In an attempt to resolve this problem I question whether the variation observed in *Centrobolus* is only because of female variation or due to a combination of male and female variation [40]. Sexual size dimorphism (SSD) is correlated with female size in the pachybolid millipede genus *Centrobolus* Cook, 1897 [5, 16, 21]. The null hypothesis is that there is an SSD correlation with female body size consistent with an inverse Rensch rule [6-11].

### 2. Materials and methods

Twenty-two valid species were identified as belonging to the genus *Centrobolus* Cook, 1897. Millipede type localities were obtained from a checklist of southern African millipedes [16]. These were tabulated and known type localities also listed in Microsoft Word online (<https://office.live.com/start/Word.aspx>) (Table 1). GPS coordinates were obtained from internet sources for known type localities using google (<https://www.google.co.za/maps/place>). Mean female size was obtained by calculating the volumes (cylindrical) using the lengths and widths of the sex which were inputted into the formula for a cylinder's volume (<https://byjus.com/volume-of-a-cylinder-calculator>) [12]. SSD was calculated as the ratio of female volume to male volume. SSD and female size was checked for correlations using the Pearson Correlation Coefficient calculator (<https://www.gigacalculator.com/calculators/correlation-coefficient-calculator.php>). Correlation coefficients between SSD and female size were compared to SSD and species size [6].

**Corresponding Author:**  
**Mark Cooper**  
 School of Animal, Plant &  
 Environmental Sciences,  
 University of the Witwatersrand,  
 Johannesburg, South Africa


### 3. Results

There was a highly significant positive correlation between SSD and female size (Fig. 1:  $r= 0.55969609$ ,  $Z$  score= $2.75652643$ ,  $n=22$ ,  $p=0.002921$ ). There was no correlation between SSD and male size. Female size was normally distributed ( $D=0.12448$ ,  $n=22$ ,  $p=0.84383$ ). SSD

was normally distributed ( $D=0.15168$ ,  $n=22$ ,  $p=0.63788$ ). The correlation coefficient between SSD and female size did not differ significantly from the correlation coefficient between SSD and species size ( $z$ -statistic= $-0.7078$ ,  $n=22,18$ ,  $p=0.4790$ ).

**Table 1:** Species in the millipede genus *Centrobolus* Cook, 1897, with SSD, type or collected localities, and female size.

Species	SSD	Location	Female size
<i>C. albitarsis</i>	2.89	Lochiel	1414
<i>C. angelicus</i>		Makhanda	
<i>C. anulatus</i>	1.19	Umhlanga Rocks	2059
<i>C. atrophus</i>		Signal Hill	
<i>C. bifidus</i>		Nkhandla	
<i>C. coriaceus</i>		cafraria	
<i>C. decoratus</i>	0.63	Ngome Forest	429
<i>C. digrammus</i>	1.01	Hout Bay	523
<i>C. dubius</i>	1.35	Gans bay	1389
<i>C. formosus</i>		cafraria	
<i>C. fulgidus</i>	1.65	Richards Bay	1888
<i>C. immaculatus</i>	2.72	Gorongosa	2309
<i>C. inscriptus</i>	1.21	Scottburgh	2245
<i>C. inyanganus</i>	1.44	Inyanga village	913
<i>C. lawrencei</i>	1.57	Pietermaritzburg	1176
<i>C. litoralis</i>		Algoa Bay	
<i>C. luctuosus</i>		Inhambambane	
<i>C. lugubris</i>	2.18	Glennconner	3491
<i>C. miniatomaculatus</i>		Tsitsikamma	
<i>C. pococki</i>		Cape Peninsula	
<i>C. promontorius</i>	0.69	Little Lions Head	231
<i>C. pusillus</i>	2.08	Qolora River mouth	1021
<i>C. richardii</i>	0.95	Richards Bay	1187.915
<i>C. ruber</i>	1.62	Port Shepstone	1850
<i>C. rubricollis</i>		Karkloof waterfall	
<i>C. rugulosus</i>	1.97	Hluhluwe	2209
<i>C. sagatinus</i>	1.27	Between Uitenhage and Addo	1855
<i>C. sanguine marginatum</i>		Bain's Kloof	
<i>C. sanguinipes</i>		Qolora River mouth	
<i>C. saussurii</i>		cafraria	
<i>C. silvanus</i>	1.13	Kentani	793
<i>C. splendidus</i>		Masiene near Chai Chai	
<i>C. strigosus</i>		cafraria	
<i>C. striolatus</i>		Port St Johns	
<i>C. titanophilus</i>	1.15	DeHoop vlei	421
<i>C. transvaalicus</i>	1.26	Mariepskop	746
<i>C. tricolor</i>	1.10	Champaigne Castle	786
<i>C. validus</i>		Haroni River	
<i>C. vastus</i>	1.81	Port St Johns	3327



**Fig 1:** Relationship between Sexual Size Dimorphism (y-axis) and female size (x-axis) in *Centrobolus* Cook, 1897 [5].

#### 4. Discussion

A relationship between female body size and SSD was found confirming the null hypothesis. *C. immaculatus* has the highest SSD (2.72) and the third-largest female size (2309 mm<sup>3</sup>). SSD was lowest in *C. promontorius* (0.69) which had the smallest female size at 231 mm<sup>3</sup> and *C. decoratus* (0.63) which had the third smallest female size (429 mm<sup>3</sup>). Only three species displayed male-biased SSD in this genus (*C. decoratus*, *C. promontorius*, *C. richardii*) and these were the exceptions to the rule of female-biased SSD among the other 19 species examined here. This study supports female size as a predictor of SSD in *Centrobolus* and helps to disentangle the size and shape components of SSD [3]. The correlation coefficient between SSD and female size did not differ significantly from the correlation coefficient between SSD and species size suggesting there was no difference between the combined effect of male and female body size [6-11] and female body size alone. Thus all the volumetric sexual size dimorphism measured in this paper and the regression with which it was compared [6] is due to variation in female width and female length.

Size-assortative mating based on female width and length is associated with variance in millipede polygynandrous mating systems across a female size gradient with higher SSD occurring in larger females [6-11]. SSD increase with female size may be explained by greater fecundity selection [6-11, 14, 32]. Female size and its components may be an explanation for skewed sex ratios in species showing sexual size dimorphisms, such as ground beetles, millipedes, scorpions, spiders, and water striders [14, 17, 20, 31-33]. Length and width components are useful reference characters because they both vary significantly with SSD and have correlated responses on female fecundity selection [8, 9].

Female-biased SSD has been successfully accepted in *Centrobolus* [6-11]. Numerous more studies have successfully rejected Rensch's rule and these include newts [1], melanine grasshoppers [2], salamanders [4], stoneflies [15], spiders [17, 20,

30], sticklebacks [18], flying lizards [19], Chinese lizards [22], frogs [23-26, 29, 40], red flour beetles [27], molluscs [28], waterstriders [31, 32], chicken breeds [33], lizards [34], ground beetles [35], dwarf chalcid wasps [36], dog breeds [37], and tinamous [38].

#### 5. Conclusion

SSD increased systematically with female size in *Centrobolus*. Morphological variance in the polygynandrous reproductive systems occurs where larger females and higher SSD occur together.

#### Competing interests

The author has declared that no competing interests exist.

#### References

1. Peñalver Alcázar M, Galán P, Aragón P. Assessing Rensch's rule in a newt: Roles of primary productivity and conspecific density in interpopulation variation of sexual size dimorphism. *Journal of Biogeography*. 2019;46:2558-2569.
2. Bidau CJ, Martí DA, Castillo ER. Rensch's rule is not verified in melanopline grasshoppers (Acrididae). *Journal of Insect Biodiversity*. 2013;1(12):1-14.
3. Cheng RC, Kuntner M. Disentangling the Size and Shape Components of Sexual Dimorphism. *Evolutionary Biology*. 2015;42:223-234.
4. Colleoni E, Denoël M, Padoa Schioppa E, Scali S, Ficetola GF. Rensch's rule and sexual dimorphism - in salamanders: patterns and potential processes. *Journal of Zoology*. 2014;293:143-151.
5. Cook OF. New relatives of *Spirobolus giganteus*. *Brandtia (A series of occasional papers on Diplopoda and other Arthropoda)*. 1897;18:73-75.
6. Cooper M. Re-assessment of Rensch's rule in *Centrobolus*. *Journal of Entomology and Zoology Studies*. 2017;5(6):2408-2410.

7. Cooper M. Allometry in *Centrobolus*. Journal of Entomology and Zoology Studies. 2018; 6(6): 284-286.
8. Cooper M. *Centrobolus* size dimorphism breaks Rensch's rule. Arthropods. 2018;7(3):48-52.
9. Cooper M. *Centrobolus* size dimorphism breaks Rensch's rule. Scholars' Press, Mauritius. 1-48, 2018. ISBN: 978-3-659-83990-0.
10. Cooper MI. Sexual size dimorphism and the rejection of Rensch's rule in Diplopoda (Arthropoda). Journal of Entomology and Zoology Studies. 2018;6(1):1582-1587.
11. Cooper MI. Trigoniuclid size dimorphism breaks Rensch. Journal of Entomology and Zoology Studies. 2018;6(3):1232-1234.
12. Cooper M. Xylophagous millipede surface area to volume ratios are size dependent in forests. Arthropods. 2019;8(4):127-136.
13. De Lisle SP, Rowe L. Correlated Evolution of Allometry and Sexual Dimorphism across Higher Taxa. The American Naturalist. 2013;182(5): 630-639.
14. Fox G, Cooper A, Hayes W. The Dilemma of Choosing a Reference Character for Measuring Sexual Size Dimorphism, Sexual Body Component Dimorphism, and Character Scaling: Cryptic Dimorphism and Allometry in the Scorpion *Hadrurus arizonensis*. PLoS ONE, 2015. 10. 10.1371/journal.pone.0120392.
15. Guillermo-Ferreira R, Novaes MC, Lecci LS, Bispo PC. Allometry for sexual size dimorphism in stoneflies defies the Rensch's rule. Neotropical Entomology. 2014;43:172.
16. Hamer ML. Checklist of Southern African millipedes (Myriapoda: Diplopoda). Annals of the Natal Museum. 1998;39(1):11-82.
17. Head G. Selection on Fecundity and Variation in the Degree of Sexual Size Dimorphism Among Spider Species (Class Araneae). Evolution. 1995;49(4):776-781.
18. Herczeg G, Gonda A, Merilä J. Rensch's rule inverted – female-driven gigantism in nine-spined stickleback *Pungitius pungitius*. Journal of Animal Ecology. 2010;79(3):581-588.
19. Husak JF, McGuire JA. Does 'gliding while gravid' explain Rensch's rule in flying lizards? Biological Journal of the Linnean Society. 2014;113:270-282.
20. Kuntner M, Coddington JA. Sexual Size Dimorphism: Evolution and Perils of Extreme Phenotypes in Spiders. Annual Review of Entomology. 2020;65:57-80.
21. Lawrence RF. The Spiroboloidea (Diplopoda) of the eastern half of Southern Africa\*. Annals of the Natal Museum. 1967;18(3):607-646.
22. Liang T, Shi L, Bempah G, Lu C-h. Sexual size dimorphism and its allometry in Chinese lizards. Evolutionary Ecology. 2021;35:323-335.
23. Liao WB. Evolution of sexual size dimorphism in a frog obeys the inverse of Rensch's rule. Evolutionary Biology. 2013;40:493-499.
24. Liao WB, Chen W. Inverse Rensch-rule in a frog with female-biased sexual size dimorphism. Naturwissenschaften. 2012;99:427-431.
25. Liao WB, Zeng Y, Zhou CQ, Jehle R. Sexual size dimorphism in anurans fails to obey Rensch's rule. Frontiers in Zoology. 2013;10(10):1-7.
26. Liao WB, Liu WC, Merilä J. Andrew meets Rensch: sexual size dimorphism and the inverse of Rensch's rule in Andrew's toad (*Bufo andrewsi*). Oecologia. 2015;177:389-399.
27. Lu D, Zhou CQ, Liao WB. Pattern of sexual size dimorphism supports the inverse Rensch's rule in two frog species. Animal Biology. 2014;64:87-95.
28. Martin OY, Michalczyk L, Millard AL, Emerson BC, Gage MJG. Lack of support for Rensch's rule in an intraspecific test using red flour beetle (*Tribolium castaneum*) populations. Insect Science. 2017;24(1):133-140.
29. Minton RL, Wang LL. Evidence of sexual shape dimorphism in *Viviparus* (Gastropoda: Viviparidae). Journal of Molluscan Studies. 2011;77(3):315-317.
30. Monnet JM, Cherry MI. Sexual size dimorphism in anurans. Proceedings of the Royal Society of London B Biological Sciences. 2002;269(1507):2301-2307.
31. Prenter J, Montgomery WI. Sexual size dimorphism and reproductive investment by female spiders: A Comparative Analysis. Evolution. 2017;53(6):1987-1994.
32. Preziosi RF, Fairbairn DJ. Lifetime selection on adult body size and components of body size in a waterstrider: opposing selection and maintenance of sexual size dimorphism. Evolution. 2000;54(2):558-566.
33. Reeve J, Fairbairn D. Change in sexual size dimorphism as a correlated response to selection on fecundity. Heredity. 1999;83:697-706.
34. Remeš V, Székely T. Domestic chickens defy Rensch's rule: sexual size dimorphism in chicken breeds. Journal of Evolutionary Biology. 2010; 23: 2754-2759.
35. Rutherford PL. Proximate mechanisms that contribute to female-biased sexual size dimorphism in an anguillid lizard. Canadian Journal of Zoology. 2004;82(5):817-822.
36. Sukhodolskaya RA, Ananina TL, Saveliev AA. Variation in Body Size and Sexual Size Dimorphism of Ground Beetle *Pterostichus montanus* Motsch. (Coleoptera, Carabidae) in Altitude Gradient. Contemporary problems of ecology. 2021;14:62-70.
37. Stuart-fox D. A test of Rensch's rule in dwarf chameleons (*Bradypodion* spp.), a group with female biased sexual size dimorphism. Evolutionary Ecology. 2009;23:425-433.
38. Sutter NB, Mosher DS, Ostrander EA. Morphometrics within dog breeds are highly reproducible and dispute Rensch's rule. Mammalian Genomics. 2008;19:713-723.
39. Tubaro PL, Bertelli S. Female-biased sexual size dimorphism in tinamous: a comparative test fails to support Rensch's rule. Biological Journal of the Linnean Society. 2003;80:519-527.
40. Webb TJ, Freckleton RP. Only half right: Species with female-biased sexual size dimorphism consistently break Rensch's rule. PLoS ONE. 2007;2(9):e897.
41. Yu TL, Li Y, Zhang JD. Evolution of sexual dimorphism in the plateau brown frog fails to obey Rensch's rule. The Herpetological Journal. 2022;32(1):27-33.