

E-ISSN: 2708-0021  
P-ISSN: 2708-0013  
[www.actajournal.com](http://www.actajournal.com)  
AEZ 2022; 3(1): 30-33  
Received: 12-11-2021  
Accepted: 24-12-2021

**Kumar Kaustubh**  
Department of Zoology, Patna  
University, Patna, Bihar,  
India

**Rahul Joshi**  
Zoological Survey of India,  
Gangetic Plains Regional  
Centre, Patna, Bihar, India

**SMM Hassan**  
Department of Zoology,  
Patna Science College, Patna  
University, Patna, Bihar,  
India

**Corresponding Author:**  
**SMM Hassan**  
Department of Zoology,  
Patna Science College, Patna  
University, Patna, Bihar,  
India

## Report of occurrence of *Lymantria mathura* (Lepidoptera: Erebidae) from Saranda Forest Division, Jharkhand, India

**Kumar Kaustubh, Rahul Joshi and SMM Hassan**

**DOI:** <https://doi.org/10.33545/27080013.2022.v3.i1a.61>

### Abstract

Saranda forest division is the famous Sal-forest of Jharkhand, India. *Shorea robusta*, commonly known as 'Sal' is an indigenous plant species, profusely regenerating and growing in the forest region. It is ecologically as well as economically important timber of India. *Lymantria mathura* Moore, [1866], commonly called Rosy Gypsy Moth (Lepidoptera, Erebidae, Lymantriinae) is an established pest species of *Shorea robusta* (Dipterocarpaceae). As a result of taxonomic identification, a major pest species of moth, *Lymantria mathura* Moore, [1866] has been identified for the first time from Saranda forest division.

**Keywords:** Saranda forest division, *Lymantria Mathura*, *Shorea robusta* and pest species

### Introduction

Saranda forest division is the famous Sal-forest of Jharkhand, India. The area shares its boundary with West Bengal and Odisha. The division comprises four forest ranges, Samta, Koina, Sasganda and Gua. It has very diverse flora which produces different types of forest products like- Sal leaves, Sal seeds, Salai gum, Siyali (*Bauhinia spp.*) leaves, Mahua (*Madhuka sp.*) flowers and fruits, Sabai grass, Honey and other plants of medicinal importance<sup>[8]</sup>. *Shorea robusta*, commonly known as 'Sal' is an indigenous plant species<sup>[7, 11]</sup>, profusely regenerating and growing in the forest region. It is ecologically as well as economically important timber of India<sup>[17, 11]</sup>. Sal trees are source of many forest products on which the local population survives.

*Lymantria mathura* Moore, [1866], the Rosy Gypsy Moth (Lepidoptera) is a polyphagous species belonging to subfamily Lymantriinae under family Erebidae. Sexual dimorphism is prominently seen in this species in terms of size and colour. Adult males have forewing brown and hindwing yellow. Females have forewings white with black markings and hindwing pink. Wingspan of adult female is found larger in comparison with the male. It is an established pest species of *Shorea robusta* (Dipterocarpaceae)<sup>[1, 2, 10, 3, 11]</sup>. Till date, there is no published report of occurrence of this moth species from Saranda Forest Division. However, Singh (2019)<sup>[14]</sup> reported *Lymantria marginata* Walker, 1855 from Saranda Forest Division.

In the present study, *Lymantria mathura* Moore, [1866] has been collected for the first time from Saranda forest division. Photograph of the adult male and female adult have been provided.

### Methodology

#### Study Area

Saranda forest division is a reserve forest situated in West Singhbhum district of Jharkhand (Figure 1). Landscape is full of hills and has rich diversity of flora and fauna. During this study, the studied species was collected from Kiriburu and B. B. Mine area (Figure 2).




Fig 1: Map of India, Jharkhand and Saranda Forest Division (Courtesy-Google India)



Fig 2: Map of Saranda Forest Division showing the localities where moths were recorded (Courtesy-Google Earth)



**Fig 3:** A survey locality (Tholkobad) in Saranda Forest Division, Jharkhand

### Collection & Identification

Adult moths were collected using vertical light traps fitted with Mercury bulb (200 W). Collection was done during pre-monsoon and post-monsoon seasons in the forest. The collected samples were treated as per recent techniques of Lepidopterology. Specimens were photographed using camera (Canon EOS 600D). All the specimens are deposited at the National Zoological Collection of Zoological Survey of India, Gangetic Plains Regional Centre, Patna, Bihar. Identification of the specimens was done with the help of relevant literature [4, 5, 12]. Detail about host plants and distribution was taken from Robinson *et al.* 2010; Shubhalaxmi *et al.*, 2011; Singh, 2019 and Singh 2022 [9, 13, 14, 15].

### Results and Discussion

**Family:** Erebidae

**Subfamily:** Lymantriinae

### Genus *Lymantria* Hübner

Hübner [1819]; *Verz.bek.Schmett.* (10): 160.

*Lymantria mathura* Moore, [1866] (Plate-1, Figure- 1 and 2)

*Lymantria mathura* Moore, 1866; *Proc. Zool. Soc. Lond.* 1865 (3): 805.

**Material Examined:** India: Jharkhand, Saranda Forest, Kiriburu (22°07.066' N; 85°17.831' E), Alt.- 897.7 m, 13.xi.2017, 1 ex-♂, 01.x.2018-1ex-♂, 02.x.2018- 2ex-♂, 03.x.2018- 1 ex-♂, 06.viii.2019-1ex.-♀, 1ex-♂, B. B. Mine (22°10.030' N; 85°21.299' E), 523.9 m asl, 08.viii.2019-1ex.-♂ (Coll.by- K. Kaustubh).


**Distribution:** India: Bihar, Jharkhand, Maharashtra, Meghalaya, North West Himalaya, Sikkim, Uttarakhand and

West Bengal (Singh, 2019). *Elsewhere:* China, Japan, Korea, Nepal (Singh, 2019).

**Host Plant:** *Lagerstroemia parviflora* (Lythraceae), *Mangifera indica* (Anacardiaceae), *Planchonia careya* (Lecythidaceae), *Quercus serrata* (Fagaceae), *Shorea robusta* (Dipterocarpaceae), *Syzygium cumini* (Myrtaceae), *Terminalia arjuna*, *Terminalia bellirica*, *Terminalia myriocarpa*, *Terminalia tomentosa* (Combretaceae) (Robinson *et al.* 2010), *Quercus griffithii* (Fagaceae) (Singh 2022 in Sondhi *et al.*).



**Plate 1:** *Lymantria mathura* Moore, [1866]



**Plate 2:** *Lymantria mathura* Moore, [1866]

## Conclusion

Since the moth fauna of this region in Jharkhand is insufficiently studied earlier, there was no published record of this species. Presence of pest species raises alarm of threat to the forest ecosystem because of dominance of a single type of plant species. Mass emergence of caterpillars of *L. mathura* caused an epidemic like situation in the Doon valley and neighbouring forest areas of Uttar Pradesh during 1953-54<sup>[10]</sup>. Caterpillars of this moth were reported to cause defoliation in Sal trees in forests of Assam and Madhya Pradesh in India<sup>[1, 3, 11]</sup>. Since, Saranda forest division is dominated by *Shorea robusta*, so it is important to know about the presence and distribution of such pest species. The moth has been recorded from the two important mining areas (Kiriburu and B. B. Mine area) of the Sasganda range of Saranda forest division, Jharkhand. Repeated surveys in the area and proper monitoring will help to ensure a check before it starts damaging like a havoc.

## Acknowledgements

Authors are thankful to the Director, Zoological Survey of India, Kolkata, Officer-In-Charge, Zoological Survey of India, G.P.R.C., Patna for providing support and encouragement during the work. We are also thankful to PCCF, Jharkhand, DFO, Saranda Forest Division, Jharkhand, Jharkhand Biodiversity Board, Jharkhand and all the forest officials and local inhabitants of the Saranda forest area of Jharkhand for their kind cooperation and help during survey tours. Last but not the least endless support and encouragement of Dr. Navneet Singh, Scientist-E & Officer-In-Charge, Lepidoptera section, Zoological Survey of India, Kolkata is profoundly acknowledged.

## Competing Interests

Authors have no conflict of interest to any extent.

## Authors' Contributions

“Author A-Kumar Kaustubh’ designed the study, performed the laboratory work, analysed the data and wrote the first draft of the manuscript. ‘Author B-Rahul Joshi’ and ‘Author C-S. M.M. Hassan’ managed the analyses of the study. All authors read and approved the final manuscript.”

## References

1. Beeson CFC. The Ecology and Control of Forest Insects of India and Neighbouring Countries. Reprint 1993. Bishen Singh Mahendra Pal Singh, Dehradun, 1941, 1007 pp.
2. Browne FG. Pests and Diseases of Forest Plantation Trees. Clarendon Press, Oxford, 1968, 1330 pp.
3. Dey RK, Tiwari KP. Detection of an imminent defoliator attack on the borer infested sal forests of Madhya Pradesh. *Vaniki Sandesh* 1997;21:21-4.
4. Hampson G. The Fauna of British India, including Ceylon and Burma. Moths, Saturniidae to Hypsiidae. Taylor and Francis, London. 1892;I:527.
5. Holloway JD. The Moths of Borneo (Lymantriidae), Part 5, The Malayan Nature Journal, Kuala Lumpur, Malaysia, 1999, 1-188.
6. James Oliver E, Joseph Dickens C, Marina Zlotinac, Victor Mastrod C, Galina Yurchenko I. Sex Attractant of the Rosy Russian Gypsy Moth (*Lymantria mathura* Moore), 1999, 387-394.

7. Kulkarni DH. Comparative distributional characteristics of sal and teak in Madhya Pradesh. Proceedings IX Silvicultural Conference, Dehra Dun, 1956.
8. Lal HS, Pramanik K, Ganguly S, Ranjan V. A Checklist of the flora of Saranda Forest Division. *Plant Systematics & Biotechnology: Challenges & opportunities*, 2018, 219-234.
9. Robinson GS, Ackery PR, Kitching IJ, Beccaloni GW, Hernández LM. HOSTS – A Database of the World’s Lepidopteran Hostplants. Natural History Museum, London, 2010. Available at-<http://www.nhm.ac.uk/hosts>. Accessed on- 05.10.2021.
10. Roonwal ML. Field-ecological studies on mass eruption, seasonal life-history, nocturnal feeding and activity rhythm, and protective behavior and coloration in the sal defoliator, *Lymantria mathura* (Lepidoptera: Lymantriidae), in sub-Himalayan forests. *Records of the Zoological Survey of India*. 1979;75:209-236.
11. Roy Choudhury N, Singh RB, Mishra RK. Occurrence of *Lymantria mathura* in sal forests of Odisha. *Van Sangyan* (ISSN 2395 - 468X). 2020;7:10.
12. Schintlmeister A. The Taxonomy of the genus *Lymantria* HÜBNER [1819] (Lepidoptera: Lymantriidae), 2004, 1- 248.
13. Shubhalaxmi V, Kendrick RC, Vaidya A, Kalagi N, Bhagwat A. Inventory of moth fauna (Lepidoptera: Heterocera) of the Northern Western Ghats, Maharashtra, India. *Journal of the Bombay Natural History Society*. 2011;108(3):183-205.
14. Singh N. Moths of Bihar and Jharkhand, *Rec. zool. Serv. India, Occasional Paper No.*, (Published by the Director, Zool. Serv. India, Kolkata). 2019;400:1-180.
15. Singh R. *Lymantria mathura* Moore, [1866] – Rosy Gypsy Moth. In Sondhi, S., Y. Sondhi, P. Roy and K. Kunte (Chief Editors). *Moths of India*, v. 2.80. Indian Foundation for Butterflies, 2022. <http://www.mothsofindia.org/sp/355457/Lymantria-mathura> - (Accessed on - 15.01.2022).
16. Singh SM. A note on serious damage to mango crop by *Lymantria mathura* Moore, in Doon Valley. *Indian Journal of Horticulture*. 1954;11:150.
17. Tewari DN. A Monograph on Sal (*Shorea robusta* Gaertn.f.). International Book Distributors, Dehradun, 1995, 277 pp.